

Pion and kaon structures from the instanton vacuum

Hyun-Chul Kim,^{1,*} Seung-il Nam,^{2,†} and Hyun-Ah Choi^{3,‡}

¹*Department of Physics, Inha University,
Incheon 402-751, Republic of Korea*

²*Yukawa Institute for Theoretical Physics (YITP),
Kyoto University, Kyoto 606-8502, Japan*

³*Department of Physics, Pusan National University,
San 30 Jangjeon-dong, Geumjung-gu, Busan, 609-735, Korea*

(Dated: September, 2008)

Abstract

In this presentation, we briefly review recent investigations on pion and kaon structures from the instanton vacuum. Starting from the low-energy QCD partition function, we have computed the Gasser-Leutwyler low-energy constants, electromagnetic form factors of the pion and kaon, semileptonic decay form factors of the kaon, and light-cone distribution amplitudes of the pion and kaon. The results are in good agreement with the experimental and empirical data.

PACS numbers: 12.38.Lg, 14.40.Ag

Keywords: Effective chiral Lagrangian, pion and kaon electromagnetic form factors, kaon semileptonic decays, light-cone distribution amplitudes, instanton vacuum.

*E-mail:hchkim@inha.ac.kr

†E-mail:sinam@yukawa.kyoto-u.ac.jp

‡E-mail:hachoi@pusan.ac.kr

I. INTRODUCTION

It is of great importance to understand structures of the pion and kaon, since they are regarded as the pseudo-Goldstone bosons arising from the spontaneous breakdown of chiral symmetry (SB χ S). Though there are many theoretical frameworks to study the pion and kaon such as chiral perturbation theory, QCD sum rules, and lattice QCD, the instanton-vacuum approach (see, for example, recent reviews [1, 2]) has certain virtues: First, it can be directly related to nonperturbative QCD, realizing well the mechanism of the SB χ S with the quark zero modes. Second, it has only two parameters: The average instanton size $\bar{\rho} \approx \frac{1}{3}$ fm and average inter-instanton distance $\bar{R} \approx 1$ fm. Thus, there is basically no free parameter in this approach. Third, the model has a natural normalization point that is given by the average size of instantons ($\rho^{-1} \approx 0.6$ GeV). It allows us to calculate hadronic observables and compare the results with the experimental data with the renormalization group employed.

The starting point is the following gauge-invariant low-energy effective chiral action:

$$\mathcal{S}_{\text{eff}}[\mathcal{M}^\alpha, V, m] = -\text{Sp}_{c,f,\gamma} \ln \left[i\bar{D} + im + i\sqrt{M(iD)}U^{\gamma 5}\sqrt{M(iD)} \right]. \quad (1)$$

One can refer to Ref. [3] for the description of Eq. (1) in detail.

II. LOW-ENERGY CONSTANTS

The SU(3) effective chiral Lagrangian to order $\mathcal{O}(p^4)$ in the large N_c and in the chiral limit is expressed as

$$\mathcal{L}^{(4)} = L_1 \langle \partial_\mu U^\dagger \partial_\mu U \rangle^2 + L_2 \langle \partial_\mu U^\dagger \partial_\nu U \rangle^2 + L_3 \langle \partial_\mu U^\dagger \partial_\mu U \partial_\nu U^\dagger \partial_\nu U \rangle, \quad (2)$$

where L_i denote the Gasser-Leutwyler low-energy constants. Utilizing the derivative expansion of Eq. (1), we can determine the L_i [4]. The results are listed in Table 1. The explanation for the abbreviation can be found in Ref. [4]. The results are in good agreement with the empirical values by Gasser and Leutwyler [5].

TABLE I: The low energy constants L_1 , L_2 , L_3 .

	$M_0(\text{MeV})$	$\Lambda(\text{MeV})$	$L_1(\times 10^{-3})$	$L_2(\times 10^{-3})$	$L_3(\times 10^{-3})$
DP	350	611.7	0.82	1.63	-3.09
Dipole	350	611.2	0.82	1.63	-2.97
Gaussian	350	627.4	0.81	1.62	-2.88
GL			0.9 ± 0.3	1.7 ± 0.7	-4.4 ± 2.5

The combination of the low-energy constant is related to the upper bound of the sigma-meson mass:

$$M_\sigma < 665[1 + 0.44\Delta + 0.33\Delta^2 + \mathcal{O}(\Delta^3)]\text{MeV}, \quad (3)$$

where $\Delta = -(2L_2 + L_3)/L_2$ which can be determined by the $\pi\pi$ scattering length. The Δ is ranged from -1.03 to -0.243, depending on the type of the form factor. As a result, the upper bound of the sigma-meson mass lies in the range of 610 ~ 640 MeV, which is very different from almost all other models in which Δ turns out to be positive.

III. ELECTROMAGNETIC FORM FACTORS OF THE PION AND KAON

The electromagnetic form (EM) factors of the pion and kaon are defined as follows:

$$\langle \mathcal{M}(P_f) | j_\mu^{\text{EM}}(0) | \mathcal{M}(P_i) \rangle = (P_i + P_f)_\mu F_{\mathcal{M}}(q^2). \quad (4)$$

The notation of Eq. (4) can be found in Ref. [3]. The results of the EM form factors of the pion and kaon are drawn in Fig. 1 (see Ref. [3] for the explanation of the abbreviation). The EM charge radii of the pion and kaon are: $\langle r^2 \rangle_{\pi^+}^{1/2} = 0.675$ fm and $\langle r^2 \rangle_{K^+}^{1/2} = 0.731$ fm




FIG. 1: The results of the electromagnetic form factors of the pion and kaon

(experimental data: $\langle r^2 \rangle_{\pi^+}^{1/2} = 0.672 \pm 0.008$ fm and $\langle r^2 \rangle_{K^+}^{1/2} = 0.560 \pm 0.031$ fm), respectively. The deviation of the kaon charge radius from the experiment is due to the fact that the $1/N_c$ corrections are absent in the present approach.

IV. SEMILEPTONIC DECAYS OF THE KAON

The corresponding matrix element of the kaon semileptonic decay form factors is written as

$$\langle \pi(p_\pi) | \bar{\psi} \gamma_\mu \lambda^{4+i5} \psi | K(p_K) \rangle = (p_K + p_\pi)_\mu f_{l+}(t) + (p_K - p_\pi)_\mu f_{l-}(t). \quad (5)$$

The detailed description for Eq. (5) can be found in Ref. [6]. The t -dependence of the form factor is shown in the left panel of Fig. 2. The results are in qualitative agreement with the data. The dimensionless slope parameter λ_{l+} is obtained as 3.03×10^{-2} (experiment: $(2.96 \pm 0.05) \times 10^{-2}$).

V. LIGHT-CONE DISTRIBUTION AMPLITUDES OF THE PION AND KAON

The gauge invariance of Eq. (1) plays an important role in deriving the light-cone distribution amplitudes (DA) of the pion [7]. In the right panel of Fig. 2, we draw the Gegenbauer moments relevant for the pion DA. The ellipses denote the Schmedding-Yakovlev analysis [8]. The results lie within the 2σ ellipsis. One can consult Ref. [7] for details.




FIG. 2: The results of the semileptonic form factors of kaon

VI. SUMMARY AND CONCLUSION

In the present talk, we briefly reviewed recent investigations on the structures of the pion and kaon from the instanton vacuum. Without adjusting any parameter, almost all results are in good agreement with the experimental data.

Acknowledgments

The authors are grateful to the organizers for the 4th Asia-Pacific Conference on Few-Body Problems in Physics 2008 (APFB08). The present work is supported by the Korea Research Foundation Grant funded by the Korean Government(MOEHRD) (KRF-2006-312-C00507). S.i.N. was partially supported by the Grant for Scientific Research (Priority Area No.17070002 and No.20028005) from the Ministry of Education, Culture, Science and Technology (MEXT) of Japan.

-
- [1] D. Diakonov, Prog. Part. Nucl. Phys. **51**, 173 (2003).
 - [2] T. Schäfer and E. V. Shuryak, Rev. Mod. Phys. **70**, 323 (1998).
 - [3] S. i. Nam and H.-Ch. Kim, Phys. Rev. D **77**, 094014 (2008).
 - [4] H. A. Choi and H.-Ch. Kim, Phys. Rev. D **69**, 054004 (2004).
 - [5] J. Gasser and H. Leutwyler, Annals Phys. **158**, 142 (1984).
 - [6] S. i. Nam and H.-Ch. Kim, Phys. Rev. D **75**, 094011 (2007).
 - [7] S. i. Nam and H.-Ch. Kim, Phys. Rev. D **74**, 076005 (2006).
 - [8] A. Schmedding and O. I. Yakovlev, Phys. Rev. D **62**, 116002 (2000).